

SYMBRAVO®

(meloxicam and rizatriptan)

20 mg/10 mg tablets

Conducting the epic party: **MOM**
Relieving migraine fast^a: **SYMBRAVO**

Bravo SYMBRAVO!

A powerful migraine treatment
to get you back to conducting the
things that matter.^a

^aIn a clinical study, some people taking SYMBRAVO experienced pain freedom and the ability to return to normal daily activities at 2 hours.

INDICATION

What is SYMBRAVO (sim-BRAH-voh)? SYMBRAVO is a combination of meloxicam (an NSAID) and rizatriptan (a triptan). SYMBRAVO is an oral prescription medicine used to treat acute migraine headaches with or without aura in adults.

SYMBRAVO is not used to prevent or decrease the number of migraine headaches you have or for treatment of hemiplegic or basilar migraines. SYMBRAVO is not indicated as a treatment for cluster headaches or for use in children.

Please see Important Safety Information throughout, including **Boxed Warning** for risk of serious cardiovascular and gastrointestinal events, and full [Prescribing Information](#).

IMPORTANT SAFETY INFORMATION

WHAT IS THE MOST IMPORTANT INFORMATION I SHOULD KNOW ABOUT SYMBRAVO?

SYMBRAVO may increase the risk of a heart attack or stroke that can lead to death. This risk may happen early in treatment and may increase with increasing doses, and longer use, of NSAIDs.

Do not take SYMBRAVO right before or after a heart surgery called a "coronary artery bypass graft" (CABG).

Avoid taking SYMBRAVO after a recent heart attack unless your healthcare provider (HCP) tells you to. You may have an increased risk of another heart attack if you take NSAIDs after a recent heart attack.



What could SYMBRAVO mean for you?

In clinical trials, SYMBRAVO showed:



Rapid pain freedom (at 2 hours vs placebo)^{a,b}



Lasting pain freedom (from 2 through 24 hours vs rizatriptan)^b



Most patients didn't use rescue medication (from 2 through 24 hours)^{a,b}



Demonstrated safety and tolerability^{a,b}

More detailed efficacy and safety information can be found on subsequent pages.

^aIn a clinical study of patients who treated early in the headache phase of a migraine attack when pain was mild.

^bIn a clinical study of patients who treated when migraine pain was moderate or severe.

IMPORTANT SAFETY INFORMATION (CONT'D)

Stop taking SYMBRAVO and get emergency help right away if you have any of the following symptoms which can be indicative of a heart attack or stroke:

- discomfort in your chest that lasts for more than a few minutes, or that goes away and comes back
- severe tightness, pain, pressure, or heaviness in your chest, throat, neck, or jaw
- pain or discomfort in your arms, back, neck, jaw, or stomach
- shortness of breath with or without chest discomfort
- breaking out in a cold sweat
- nausea or vomiting
- feeling lightheaded
- weakness in one part or one side of your body
- slurred speech

People with risk factors for heart disease should not take SYMBRAVO unless a heart exam is done and shows no problem. You have higher risk for heart disease if you:

- have high blood pressure
- have high cholesterol
- smoke
- are overweight
- have diabetes or a family history of diabetes

SYMBRAVO can increase the risk of potentially life-threatening bleeding, ulcers, and tears (perforation) of the esophagus (tube leading from the mouth to the stomach), stomach, and intestines that can occur anytime during use and without warning symptoms.



Harnessing MoSEIC™ rapid-absorption technology to create a novel migraine medication

No combination of generic meloxicam and rizatriptan will result in the same amounts of drug reaching the bloodstream as SYMBRAVO because of its formulation

IMPORTANT SAFETY INFORMATION (CONT'D)

SYMBRAVO may cause serious allergic or skin reactions which can be life-threatening. Stop taking SYMBRAVO and get emergency help right away if you develop:

- sudden wheezing or problems breathing or swallowing
- rash or reddening of your skin with blisters or peeling
- blisters or bleeding of your lips, eye lids, mouth, nose, or genitals
- swelling of your lips, tongue, throat or body
- fainting

SYMBRAVO already contains an NSAID (meloxicam). Do not use SYMBRAVO with other medicines to lessen pain or fever or with other medicines for colds or sleeping problems without talking to your HCP first, because they may contain an NSAID also.

Do not take SYMBRAVO if you:

- have or had heart problems or right before or after heart bypass surgery
- have or had a stroke or transient ischemic attack (TIA)
- have or had blood vessel problems of your legs and arms, stomach (ischemic bowel disease), or kidneys
- have or had hemiplegic or basilar migraines
- have uncontrolled high blood pressure
- take propranolol containing medicines

- have taken other triptan or ergot-containing medicines within the last 24 hours
- take an antidepressant medicine called monoamine oxidase inhibitor (MAOI) or have taken a MAOI within the last 2 weeks
- are allergic to meloxicam, rizatriptan, NSAIDs, or any of the ingredients in SYMBRAVO
- have had an asthma attack, hives, or other allergic reaction after taking aspirin or any other NSAIDs
- have moderate to severe kidney problems and are at risk of kidney failure or if you are on dialysis

SYMBRAVO may cause serious side effects. These serious side effects include:


- heartbeats that are too fast or too slow (arrhythmias)
- new or worse high blood pressure
- heart failure
- life-threatening skin reactions
- liver or kidney problems including organ failure
- low red blood cell count (anemia)
- asthma attacks in people who have asthma

Please see Important Safety Information throughout, including **Boxed Warning** for risk of serious cardiovascular and gastrointestinal events, and full [Prescribing Information](#).

SYMBRAVO®
(meloxicam and rizatriptan)
20 mg/10 mg tablets

SYMBRAVO is thought to work differently

Migraine is a complex condition with many factors that contribute to symptoms during an attack.



The exact mechanisms causing migraine and the exact way SYMBRAVO works are not fully understood.

CGRP, calcitonin gene-related peptide; PGE, prostaglandin E.

IMPORTANT SAFETY INFORMATION (CONT'D)

Medication Overuse Headaches: Some people who use too many SYMBRAVO tablets may have worse headaches. If your headaches get worse, your HCP may decide to stop your treatment with SYMBRAVO.

Stop taking SYMBRAVO and get emergency help right away if you have any of the following:


- **Stomach and intestinal problems.** Symptoms of gastrointestinal and colonic ischemic events may include sudden or severe

stomach pains even after meals; sudden weight loss; severe nausea, vomiting, constipation, diarrhea; and bloody diarrhea.

- **Circulation problems to legs and feet.** Symptoms of peripheral vascular ischemia may include cramping and pain in your legs and hips; heaviness or tightness in leg muscles; burning, aching, numbness, tingling, or weakness in your legs, feet, or toes; cold feelings or color changes in one or both legs or feet.

Harnessing rapid-absorption technology

SYMBRAVO combines rizatriptan, one of the most prescribed migraine treatments, with long-lasting^a meloxicam and MoSEIC™ technology to create a novel migraine medication.



^aLong-lasting refers to duration of medication in blood. Blood levels do not necessarily tie to efficacy.

The exact way that SYMBRAVO works to treat migraine is unclear, and whether MoSEIC™ (Molecular Solubility Enhanced Inclusion Complex) has an impact on the effectiveness of SYMBRAVO has not been established. The reason for the faster absorption seen with rizatriptan in SYMBRAVO is unknown.

Please see Important Safety Information throughout, including **Boxed Warning** for risk of serious cardiovascular and gastrointestinal events, and full [Prescribing Information](#).

SYMBRAVO®
(meloxicam and rizatriptan)
20 mg/10 mg tablets

Powerful performance

If you aren't getting the migraine pain relief

In one clinical study,^a people took SYMBRAVO early in their migraine pain when it was still mild.



RAPID PAIN FREEDOM AT 2 HOURS

Of people who took SYMBRAVO:

33% achieved pain freedom vs placebo (16%)

44% achieved freedom from most bothersome symptom^b vs placebo (27%)



LASTING PAIN FREEDOM, 2 THROUGH 24 HOURS

Of people who took SYMBRAVO:

23% achieved lasting pain freedom vs placebo (13%)

70% of those who achieved pain freedom at 2 hours did not experience a headache return

85% did not use rescue medication vs placebo (58%)

^aIn a single-dose study of 132 patients treated with SYMBRAVO and 135 treated with placebo.

^bNausea, light sensitivity, or sound sensitivity.

IMPORTANT SAFETY INFORMATION (CONT'D)

- **Serotonin syndrome.** Can happen when taking SYMBRAVO with antidepressant medicines called SSRIs or SNRIs. Stop taking SYMBRAVO and call your doctor right away if you have any of the following symptoms:
 - mental status changes including agitation, hallucinations, or coma
 - fast heartbeat
 - changes in your blood pressure

worth cheering about

that you need, SYMBRAVO may help

In another clinical study,^a people took SYMBRAVO when their pain had progressed to moderate to severe. The study focused on people who had not achieved adequate response from their prior migraine medicine.



RAPID PAIN FREEDOM AT 2 HOURS

Of people who took SYMBRAVO:

20% achieved pain freedom vs placebo (7%)

37% achieved freedom from most bothersome symptom^b vs placebo (24%)

~70% of patients achieved pain relief^c vs placebo (55%)



LASTING PAIN FREEDOM, 2 THROUGH 24 HOURS

Of people who took SYMBRAVO:

16% achieved lasting pain freedom vs rizatriptan (11%)

81% of those who achieved pain freedom at 2 hours did not experience a headache return

77% did not use rescue medication vs rizatriptan (65%)

^aIn a single-dose study of patients treated with SYMBRAVO (428 patients); rizatriptan (419); MoSEIC™ meloxicam (421); and placebo (209).

^bNausea, light sensitivity, or sound sensitivity.

^cPain relief was defined as a reduction in migraine pain intensity from moderate or severe to mild or no headache pain.

IMPORTANT SAFETY INFORMATION (CONT'D)

- increased body temperature
- tight muscles
- trouble walking

Please see Important Safety Information throughout, including **Boxed Warning** for risk of serious cardiovascular and gastrointestinal events, and full [Prescribing Information](#).

SYMBRAVO®
(meloxicam and rizatriptan)
20 mg/10 mg tablets

Demonstrated safety and tolerability



In clinical studies, SYMBRAVO was well tolerated. The most common side effects were dizziness (2%) and tiredness (2%) in single-dose migraine treatment studies.



During a study of over 700 patients that lasted up to 12 months with ~19,000 doses of SYMBRAVO, 2% of patients left due to side effects.



IMPORTANT SAFETY INFORMATION (CONT'D)

Stop taking SYMBRAVO and call your healthcare provider right away if you have any of the following symptoms:

- nausea
- vomiting blood
- more tired or weaker than usual
- blood in your bowel movement or it is black and sticky like tar
- diarrhea
- itching, skin rash, or blisters with fever
- unusual weight gain
- your skin or eyes look yellow
- indigestion or stomach pain

8

Taking SYMBRAVO

- Shown to be effective whether taken when migraine pain was mild or moderate to severe. Encouraged to use at the earliest sign of headache pain
- Clinical trials included some patients taking a preventative migraine medicine
- Can be taken with or without food
- Maximum daily dose is 1 pill as needed
- Take as instructed by your healthcare provider
- Do not use with other NSAIDs without talking to your healthcare provider

You can find more information about taking SYMBRAVO in the full [Prescribing Information](#).

Talk to your doctor to learn more about SYMBRAVO, and visit www.SYMBRAVO.com

- swelling of the arms, legs, hands, or feet
- flu-like symptoms
- tenderness in your right upper side
- vision problems

COMMON SIDE EFFECTS

The most common side effects of SYMBRAVO include dizziness and tiredness.

These are not all the possible side effects of SYMBRAVO. Tell your doctor if you have any side effects. You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

IMPORTANT SAFETY INFORMATION (CONT'D)

BEFORE USING

- Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.
- It is important to tell your HCP if you are taking:
 - Propranolol containing medicines such as Inderal® LA or Innopran® XL
 - Aspirin or other anti-coagulants (blood thinners)
 - Medicines to help your mood including SSRIs and SNRIs
- If you are unsure if you take any of these medicines, ask your HCP. They can tell you if it is safe to take SYMBRAVO with your other medicines.
- Tell your HCP if you are pregnant or plan to become pregnant. SYMBRAVO is not recommended during pregnancy. Taking NSAIDs, including SYMBRAVO, at about 20 weeks of pregnancy or later may harm your unborn baby. NSAIDs, including SYMBRAVO, should not be taken after about 30 weeks of pregnancy.
- Tell your HCP if you are breastfeeding or plan to breastfeed.

Tell your HCP about all your medical conditions, including if you:

- have or have had heart problems, high blood pressure, chest pain, or shortness of breath
- have any risk factors for heart or blood vessel problems
- have kidney or liver problems
- have asthma

Review the list below with your HCP. SYMBRAVO may not be right for you if:

- take daily preventative aspirin
- you are pregnant or plan to become pregnant
- you are breastfeeding or plan to breastfeed

Please see Important Safety Information throughout, including **Boxed Warning** for risk of serious cardiovascular and gastrointestinal events, and full [Prescribing Information](#).

HOW TO TAKE

- SYMBRAVO is available by prescription only.
- Take SYMBRAVO exactly as instructed by your HCP.
- The maximum daily dose of SYMBRAVO is 1 tablet. Talk to your HCP about what to do if your headache does not go away or comes back.
- Take SYMBRAVO for the shortest time needed.
- Swallow SYMBRAVO tablets whole. Do not crush, chew, or divide the tablets.
- SYMBRAVO can be taken with or without food.
- Do not give SYMBRAVO to other people.
- If you take too much SYMBRAVO call your poison control center at 1-800-222-1222 or go to the nearest hospital emergency room right away.


LEARN MORE

For more information about SYMBRAVO, call 866-496-2976 or visit SYMBRAVO.com.

This summary provides basic information about SYMBRAVO but does not include all information known about this medicine. Read the information that comes with your prescription each time your prescription is filled. This information does not take the place of talking with your doctor. Be sure to talk to your doctor or other HCP about SYMBRAVO and how to take it. Your HCP is the best person to help you decide if SYMBRAVO is right for you.

SYM CON BS 01/2025

SYMBRAVO®
(meloxicam and rizatriptan)
20 mg/10 mg tablets



SYMBRAVO®
(meloxicam and rizatriptan)
20 mg/10 mg tablets



As little as \$0* per month

With the SYMBRAVO Savings Card you could pay as little as \$0 for your monthly prescription. Use the QR code or visit www.SYMBRAVO.com/SaveNow

*For patients with commercial insurance only. Offer not available to patients with Medicare, Medicaid or other government-funded insurance.

Learn more at www.SYMBRAVO.com

Please see Important Safety Information throughout, including **Boxed Warning** for risk of serious cardiovascular and gastrointestinal events, and full [Prescribing Information](#).

SYMBRAVO, AXSOME, and their logos are trademarks or registered trademarks of Axsome Therapeutics, Inc. or its affiliates. Other trademarks are property of their respective owners.

© 2025 Axsome Therapeutics, Inc. All rights reserved. PP-SYM-US-2500060 06/25

axsome

